

Evaluation of the Effects of Potassium Fertilizer on Potato Growth and Yield in Saboti Sub County, Trans Nzoia County, Kenya

MARTIN SANGULA NGOMAT

DR. MUGAMBI MWORIA

MASTER OF SCIENCE IN AGRICULTURAL AND RURAL DEVELOPMENT OF KENYA METHODIST UNIVERSITY

Email: martinngomat@yahoo.com

Abstract Irish potatoes are an important field crop grown for food. Successful Irish potato production is often hindered by lack of fertilizer application and poor choice of Potato planting varieties and hence low yield. The purpose of the study was to assess potato growth and yield response to application of potassium chlorid(muriate of potash) in Milimani Location, Saboti Sub-county, Trans Nzoia county. A randomized complete block design experiment with three replicates was conducted. Growth and yield data collected was subjected to Analysis of Variance (ANOVA) and means separated by Duncan Multiple Range Test (DMRT) at $p < 0.05$ Significance level using SAS software. It was found out that there was significant difference in yield and yield attributes. However there were no significant differences in mean yield between the two sites of production. Fertilizer rates differed significantly. It was recommended that the fertilizer industry should change fertilizer formulation; establish fertilizer blending plants and the ministry of agriculture to establish a fertilizer advisory service for farmers to boost harvest and incomes and that the fertilizer rate of 250kg/ha be applied and Sherekea variety be variety of choice for Milimani Location, Saboti Sub-county, Trans Nzoia County.

Key words Irish, Potatoes, Fertilizer, Variety, Potassium chloride, Rate

DOI: 10.6007/IJAREG/v4-i1/3737

URL: <http://dx.doi.org/10.6007/IJAREG/v4-i1/3737>

1. Introduction

Potato is one of the most consumed produce in Kenya, second to maize. Potato is grown in Kenya by approximately 500,000 small scale farmers on 120,000 hectares with yield of 7.7 tons/ha (Janssens, 2013). The rise in potato production is as a result of an increase in total area under cultivation. Productivity has fallen by 17.4% between 2010 and 2013 (FAOSTAT 2014). Despite increased acreage in production of potatoes, statistics indicate that yield per acre is reducing and this implies that certain variables are responsible causes. According to Rytel et al. (2013), the quality of potato tubers and their chemical composition are influenced by genetics, soil fertility, weather conditions and chemical treatments that are applied. Therefore, additional K-containing sources are necessary to supply to provide optimal plant growth performance and yield (Zorb et al., 2014). Rhue et al., (1986) reported that maintaining optimal K nutrition is of a big importance for development of potato plants. Potassium influences synthesis, location, transformation and storage of carbohydrates, tuber quality and processing characteristics as well as plant resistance to stress and diseases (Ebert, 2009). According to Janssens (2013), production

of potatoes in Kenya is characterized by a couple of constrains. The first initial constraint is that production is bimodal. It is produced twice a year following rainfall pattern of Kenya. Other constraints include diseases like Late Blight and Brown Rot, Lack of crop rotation where farmers cultivate potatoes in the same piece of land over and over again, poor storage facilities and lack of enough capital for capital intensive production which increases production levels.

1.1. Research Questions

1. To assess the relationship between potassium chloride application rate to the growth of potatoes.
2. To evaluate the relationship between potassium chloride application rate to the weight of different sizes of the three potato varieties.
3. To compare the relationship between potassium chloride application rates to the yield of three potato varieties.

1.2 Research Objectives

1. There is no significant relationship in the application of potassium chloride rates and the growth parameters of potato varieties
2. There is no significant difference in the application of potassium chloride application rates to the weight and size of three Irish potato varieties.
3. Application of potassium chloride fertilizer does not significantly affect the yields of three potato varieties

2. Literature review

2.2 Potassium in the soil and availability to plants

Potassium application has been neglected by majority of farmers in our country resulting in continual depletion of soil K (Ladha et al., 2003 and Lai et al, 2007) Indian soils are generally high in total K but only a small fraction of it is present in available form because of dynamic equilibriums between exchangeable, non exchangeable and fixed K With high crop intensity and high K removal, the soils are becoming deficient in K. Frequent K deficiencies have been observed in crops in this region (Panullah et al, 2006). Regmi et al. (2002) suggested that because of inadequate K application, soil K imbalance in agricultural ecosystem and stagnation of yield will become more pronounced with time. The major constraint to potato production in the cool highlands of Kenya is the rapid decline in soil fertility occasioned by continuous cultivation without adequate replenishment of mined nutrients (Kiiya et al., 2006). The situation is exacerbated by the inherently high soil acidity with pH values of 4 to 5 being common (Recke, 1997; Kiiya et al., 2006). Due to small land sizes, farmers continuously plant crops on the same land, practicing intensive cropping systems that mainly involve double and relay cropping of different crops without a fallow period (Kaguongo et al, 2008). Fertilizer is mostly applied below the recommended rate of (90 kgN/ha + 230 P₂O₅/ha) (Kaguongo et al., 2008).

2.2. Potassium chloride application rates to the weight and size of three Irish potatoes


Potatoes remove very large amounts of K in the harvested tubers (200 to 400 lb/A), but K is taken up by the plant in the cationic K⁺ form, regardless of the fertilizer material applied to the soil. Potassium deficiencies reduce the yield, size, and quality of the potato crop. A lack of adequate K can be associated with low specific gravity in potatoes. During peak periods, potatoes can take up 14 lb K/A each day. Excessively high K can result in its accumulation in tubers, where it -may increase the water content and decrease the specific gravity. Potassium fertilizer is commonly broadcast across the field prior to planting. Band placement of a small fraction of K fertilizer in the beds or during row mark out will also efficiently supply K to the plants. Since potatoes have a very high K requirement, it is important that an adequate

supply be maintained in the root zone at all times. Many important yield and quality parameters of the tuber are negatively affected if the K supply runs short, especially during tuber bulking. The harvested tubers may contain over 90% of the total K taken up by the plant. Pre -plant applications are the most effective way to supply K. Avoid high rates of banded K in close proximity to the seed piece. A combination of broadcast and band Reference placement may be more appropriate if K application rates are high. Broadcast application of K shortly after plant emergence is also common.

2.3. Effect potassium source on potato tuber quality

Potato plants absorb large quantities of potassium throughout the growing season and are known as a potassium preferring crop (Kelling et al. 2002; Blagoeva et. al., 2004). By exploring commercial potato fields, Dow et al. (1978) established that potassium was allocated from fields to potato tubers in the range from 203 to 397 lb K/acre which was directly proportional to tuber yields. Therefore, relatively high levels of potassium fertilization are usually recommended to achieve optimal potato yield. Positive influence of increasing potassium rates on potato yield was established by Imas and Bansal (1999); Kelling et al. (2002) and Al-Moshileh et al. (2005). Studies have shown that increasing rates of K resulted in increased tuber formation as the maximum yield was achieved at K400 and K600 when fertilized with K₂SO₄ and KCl respectively. results were obtained by Kelling et. al. (2002) who found no significant and distinct influence of K source on potato yield. In contrast to positive influence of potassium fertilization rate on potato tuber yield, the effect of the source of potassium was not found significant by Stanley and Jewell (1989) as well. However Qin et. al. (2008) did not observe yield differences due to the source of potassium. Dry matter content is an essential quality parameter for potato processing. High dry matter content improves the quality of potatoes designated for human nutrition (Wibowo et. al., 2014). It corresponds to high chips yield and low oil absorption during frying. Furthermore, reduced dry matter content in tubers makes them more susceptible to mechanical damage (Ishpekov et. al. 2000). The observed effect was more pronounced, in treatments with KCl where the decrease of dry matter content reached 15% at the highest K level (K600) when compared to the control. Similar observation was reported by Bansal and Trehan (2011) who established enhanced reduction of dry mater content in tubers when fertilized with KCl. The effect of KCl on starch content in potato tubers was even more extreme.

Conceptual Framework



3. Methodology of research

3.1. Research Design

The study used was a randomized complete block design (RCBD), Number of replications were three. Total number of plots per site was 36 with two sites which gave a total number of plots – 72 .Plot sizes 2m x 2m= (4m²).Total field area – 30m x 8 m – 240 m². Alleyway of - 1m left between blocks.

3.2. Data Collection Instruments and Sampling Techniques

Data was collected by observing and measuring different experimental units. Among were: the number of stems per plant, Stand count, Plant height, Number of leaves per plant, Aggregate and grade wise tuber yield and Leaf Area Index (LAI)

3.3. Data Analysis

The data from the measured variables was subject to analysis of variance (ANOVA) to determine whether there was a significant difference between the variable means. The means were separated by Lsd at (P≤0.05) significance level using SAS programme version 9.1

3.4. Results of the Study

4.1 Fertilizer rate and variety on growth and yield of Irish potatoes

Fertilizer rate and variety on Leaf Area Index of Irish potatoes

In terms of leaf area index (L.A.I) there was a significant difference between different fertilizer rates. 0 kg/ha produced plants with smaller L.A.I. The L.A.I increased with the increase in fertilize rate. Varieties Sherekea and markies were better in terms of L.A.I as compared to Kenya Mpya in both Meru and Milimani sites.(fig2)




Figure 1: Fertilizer rate and variety on Leaf Area Index of Irish potatoes

Fertilizer rate and variety on weight of small size of Irish potatoes

There was a significant difference (P<0.03) difference between small and cocktail sized potatoes. Generally the size increased with increase in fertilizer rate with 150kg/ha to giving the heaviest potatoes and then started to decreased at 250kg/ha. Varieties Sherekea and markies produced the heaviest potatoes while Kenya Mpya produced the lightest (Fig 7c, and 7 d)




Figure 7c: Fertilizer rate and variety on weight of small size of Irish potatoes

4.3 Fertilizer rate and variety on yield of Irish potatoes

Yield showed that there was a significant difference ($P < 0.05$) between fertilizer level and variety. The yield increased with increase in fertilizer rate. Generally 150kg/ha produced the highest yield. Sherekea did better by producing an average of 8tons/ ha. Both Meru and Milimani were not significantly different ($P < 0.05$) in terms of yield. Fig 6.




Figure 2: Fertilizer rate and variety on yield of Irish potatoes

Conclusions and discussions

It was concluded from the study that, Potassium chloride application improves the growth of Irish potatoes. There were indeed significant differences in varietal tuber yield with the application of potassium chloride, with variety Sherekea producing the highest yield of 10% advantage over markies and Kenya Mpya varieties respectively. Application of potassium chloride influenced the weight in different sizes as it increased the proportion of large tubers relative to small ones. Application rate of potassium chloride at 250kg/ha for Milimani location gave optimum production level of potatoes. The fertilizer used as a side dressing/ top dressing.

5.3 Recommendations

From the study findings it was recommended that, application of a compound fertilizer rich in K is recommended for increased yield in production of potatoes relative to fertilizer with less or none of this nutrient like DAP. It was recommended fertilizer formulation specific to meet potato production requirements for each specified agro-zones. Establishment of fertilizer blending plants and fertilizer advisory service for farmers to boost potato production and incomes. The fertilizer rate of 250kg/ha and variety Sherekea is recommended for Milimani Location Saboti Sub-county.

5.4 Suggestion for further research.

Research context was limited to Saboti in Trans Nzoia west Sub County, therefore the future research can be extended to different sectors of sub counties and counties.

References

1. Abay, A. and Sheleme, B. (2011) The Influence of Potassium Fertilizer on the Production of Potato (*Solanum tuberosum*L.) at Kembata in Southern Ethiopia. *Journal of Biology, Agriculture and Healthcare*, 1, 1-12.
2. Abd El-Latif, K.M., Osman, E.A.M., Abdullah, R. and Abdel Kader, N. (2011) Response of Potato Plants to Potassium
3. Abong, G.O., Okoth, M.W., Imungi, J.K. and Kabira, J.N. (2010) Evaluation of Selected Kenyan Potato Cultivars for Processing into Potato Crisps. *Agriculture and Biology Journal of North America*, 1, 886-893.
4. Abubaker, S., Abu Rayyan, A., Amre, A., Alzu'bil, Y. and Hadidi, N. (2011) Impact of Cultivar and Growing Season on Potato under Center Pivot Irrigation System. *World Journal of Agricultural Sciences*, 7, 718-721.
5. Adhikary, B.H. and Karki, K.B. (2006) Effect of Potassium on Potato Tuber Production in Acid Soils of Malepatan, Pokhara. *Nepal Agriculture Research Journal*, 7, 42-48.
6. Agriculture, Haile Sellasie I University, Experiment Station, Dire Dawa, Ethiopia, No 44.
7. Ali, M.R., Costa, D.J., Abedin, M.J., Sayed, M.A. and Basak, N.C. (2009) Effect of Fertilizer and Variety on the Yield of Sweet Potato. *Bangladesh Journal of Agricultural Research*, 34, 473-480.
8. Al-Moshileh, A.M. and Errebi, M.A. (2004) Effect of Various Potassium Sulfate Rates on Growth, Yield and Quality of Potato Grown under Sandy Soil and Arid Conditions. IPI Regional Workshop on Potassium and Fertigation Development in West Asia and North Africa, Rabat, 24-28 November 2004. and *Rural Development*, 3, 296-310.
9. Asmaa, R. M., and Magda M. H. Increasing productivity of potato plants (*solanum*
10. Asmaa, R.M. and Magda, M.H. (2010) Increasing Productivity of Potato Plants by Using Potassium Fertilizer and
11. Awad, S. M. The influence of organic and mineral fertilization on growth, yield and quality
12. Bansal, S.K. and Trehan, S.P. (2011) Effect of Potassium on Yield and Processing Quality Attributes of Potato. *Karnataka Journal of Agricultural Sciences*, 24, 48-54.
13. Bergmann, W. (1992) *Nutritional Disorders of Plants Development, Visual and Analytic Diagnosis*. Gustav Fischer, Jena.
14. Biniam, M.G., Githiri, S.M., Tadesse, M. and Remmy, W.K. (2014) Diagnostic Survey on Potato Production Practices in Eritrea. *ARPN Journal of Agricultural and Biological Science*, 9, 444-453

15. Biniam, M.G., Githiri, S.M., Tadesse, M. and Remmy, W.K. (2014b) Potato Seed Supply, Marketing and Production Constraints in Eritrea. *American Journal of Plant Sciences*, 5, 3684-3693. <http://dx.doi.org/10.4236/ajps.2014.524384>
16. Biniam, M.G., Githiri, S.M., Tadesse, M. and Remmy, W.K. (2016) Morphological Diversity of Farmers' and Improved Potato (*Solanum tuberosum*) Cultivars Growing in Eritrea. *Journal of Plant Studies*, 5, 63-71.
17. Blakemore, M. 1966. Seasonal changes in the amounts of phosphorus and potassium dissolved from soils by dilute calcium chloride solutions. *Journal of Agricultural Science, Cambridge* 66:139–146. 05–812
18. Brady, N.C. and Weil, R.R. (2002) *The Nature and Properties of Soil*. 13th Edition, Prentice Hall, Upper Saddle River, New Jersey.
19. Bray, R.H. 1944. Soil-plant relations I. The quantitative relation of exchangeable potassium to crop yields and to crop response to potash additions. *Soil Science* 58:305–324.
20. Cultivars under different fertilization levels in sandy soils. 1999, Ph.D. Thesis Fac. of Agric.
21. Dampney, P., Wale, S. and Sinclair, A. (2011) Review Potash Requirements of Potatoes. Report of Agriculture & Horticulture Development Board 2011.
22. Enujeke, E.C. (2013) Effects of Variety and Spacing on Growth Characters of Hybrid Maize. *Asian Journal of Agriculture*
23. FAO (2008) *Production Year Book*. Food and Agriculture Organization of the United Nations, Rome.
24. FAO (2009) Potato and Food Price Inflation. International Year of the Potato 2008. Food and Agriculture Organization Fertilizer Rates and Soil Moisture Deficit. *Advances in Applied Science Research*, 2, 388- 397.
25. Gildemacher, P.R. (2012) Innovation in Seed Potato Systems in Eastern Africa. PhD Thesis, Wageningen University,
26. Gunasena, H.P.M. and Harris, P.M. (1971) The Effect of CCC, Nitrogen and Potassium on Growth and Yield on Two Varieties of Potato. *The Journal of Agricultural Science*, 76, 33-52.
27. Haile, W. (2009) On Farm Verification of Potassium Fertilize Effect on the Yield of Irish Potato Grown on Acidic
28. Havlin, J.L., Beaton, J.D., Tisdale, S.L. and Nelson, W.L. (2005) *Soil Fertility and Fertilizers: An Introduction to Nutrient Management*. 7th Edition, Pearson Educational, Inc., Upper Saddle River, New Jersey.
29. Hoskinson, R.L., Hess, J.R., and Alessi, R.S. 1999. Temporal changes in the spatial variability of soil nutrients. In J.V. Stafford (ed.). *Precision Agriculture '99*. Part 1. Academic Press, Sheffield, England. p. 61–70.
30. Humic Acid Application. *International Journal of Academic Research*, 2, 83-88.
31. Institute, CH the Hague. www.potato.nlInstitute, Halhale, Eritrea.
32. Iritani, W.M., Weller, L.D. and Knowles, N.R. (1983) Relationship between Stem Numbers, Tuber Set and Yields of Russet Burbank Potatoes. *American Potato Journal*, 60, 423-431. <http://dx.doi.org/10.1007/BF02877248>
33. Kay, R.D., Edwards, W.M. and Duffy, P.A. (2012) *Farm Management*. 7th Edition, McGraw-Hill Companies, Inc.,
34. Luebs, R.E., Stanford, G., and Scott, A.D. 1956. Relation of available potassium to soil moisture. *Soil Science Society of America Proceedings* 20:45–50.
35. Marschner, H. (1995) *Mineral Nutrition of Higher Plants*. 2nd Edition, Academic Press Limited, San Diego.

36. Ministry of Agriculture (2010) Vegetable Crops Research Program. Annual Report of National Agricultural Research
37. Ministry of Agriculture (2011) Annual Report. Horticulture Division Report, Asmara, Eritrea.
38. Muriithi, M.M. and Irungu, J.W. (2004) Effect of Integrated Use of Inorganic fertilizer and Organic Manures on Bacterial
39. Murphy, H.F. (1968) A Report on the Fertility Status and Other Data on Some Soils of Ethiopia. Bulletin, College of New York.
40. NIVAP (2011) Netherland Catalogue of Potato Variety. AC Den Haag, The Netherlands. www.nivap.nl
41. Noor, M.A. (2010) Physiomorphological Determination of Potato Crop Regulated by Potassium Management. Ph.D. of potato crop .J. Agric. Sci. Mansoura Univ. 2005, 30: 7965-7975. of the United Nations, Rome.
42. Peirce, L.C. (1987) Vegetables: Characteristics, Production and Marketing. John Wiley and Sons, Inc., Hoboken.
43. Pervez, M.A., Ayyub, C.M., Shabeen, M.R. and Noor, M.A. (2013) Determination of Physiomorphological Characteristics of Potato Crop Regulated by Potassium Management. Pakistan Journal of Agricultural Sciences, 50, 611-615.
44. Rana, M.K. (2008) Olericulture in India. Kalyani Publishers, Ludhiana. Res., 2. 83-88.
45. Sajjan, A.S., Shekhargounda, M. and Badanur (2002) Influence of Data of Sowing, Spacing and Levels of Nitrogen on Yield Attributes and Seed Yield of Okra. Ikamataka. *Journal of Agricultural Science*, 15, 267-274.
46. Soils of Hagere Selam, Southern Ethiopia. Ethiopian Journal of Natural Resources, 11, 207-222.
47. Tadesse, M. (2000) Manipulating the Physiological Quality of in Vitro Plantlets and Transplants of Potato. PhD Thesis, submitted to Wageningen University, The Netherlands.
48. Thesis, Institute of Horticultural Sciences University of Agriculture, Faisalabad.
49. Tindall, T. and Westermann, D.T. (1994) Potassium Fertility Management of Potatoes. University of Idaho Potato School (Mimeo). Idaho State University, Pocatello. tuberosum, I.) By using potassium fertilizer and humic acid application, 2010, I. J. Academic
50. Vaezzadeh, M. and Naderidarbaghshahi, M. (2012) The Effect of Various Nitrogen Fertilizer Amounts on Yield and Nitrate Accumulation in Tubers of Two Potato Cultivars in Cold Regions of Isfahan (Iran). *International Journal of Agriculture and Crop Sciences*, 4, 1688-1691.
51. Van der Zaag, P. (1981) Soil Fertility Requirements for Potato Production. Technical Information Bulletin 14, CIP, Lima.
52. Van der Zaag, W.D.E. (1992) Potatoes and Their Production in the Netherlands. 3rd Edition, Netherlands Potato Consultative Wageningen, The Netherlands.
53. Wibowo, C., Wijaya, K., Sumartono, G.H. and Pawelzik, E. (2014) Effect of Potassium Level on Quality Traits of Indonesian Potato Tubers. *Asia Pacific Journal of Sustainable Agriculture Food and Energy*, 2, 11-16.
54. Wilt Incidence (BWI) and Tuber Yield in Potato Production Systems on Hill Slopes of Central Kenya.